

IMMUNIZATION SIDE EFFECTS

DTaP/Tdap	Diphtheria, Tetanus, Acellular pertussis
IPV	Injectable Polio
Pentacel	DTaP/Hib/IPV
Prenvar	Pneumococcal
Menactra	Meningococcal
	<ol style="list-style-type: none">1. Fever may appear up to 48 hours, if longer and over 102 rectally, call the doctor at the office.2. Fever can be as high as 102 F or 38.8 C rectally3. The injection site may be sore, swollen, or red transiently. Cold compresses may be applied. With Tetanus, a lump can last 1-2 months.4. Irritability may occur within 6-12 hours after the injection and may last for 24 hours.5. Sleepiness can also occur on the day of the vaccines.
Hep A	Hepatitis A
Hep B	Hepatitis B
Hib	Haemophilus Influenza B
Gardasil	Human Papillomavirus (HPV)
	<ol style="list-style-type: none">1. Fever may appear up to 48 hours, if longer and over 102 F rectally, call the doctor at the office.
MMR	Measles, Mumps, Rubella
	<ol style="list-style-type: none">1. Reaction may occur 7-21 days after vaccination.<ol style="list-style-type: none">a. Low grade fever.b. Light pinpoint rash all over body.c. If reaction occurs, it is not communicable.
Varivax	Varicella Virus (Chickenpox)
	<ol style="list-style-type: none">1. Reaction may occur 5-26 days after vaccination<ol style="list-style-type: none">a. Low grade fever.b. 5-10 pox lesions (look like mosquito bites).c. Blisters at the injection site.d. Contagious only if rash is present.e. Avoid newborns, pregnant women who have not had chicken pox and anyone on chemotherapy or immunocompromised.2. There is no guarantee of perfect lifelong immunity; however, Vaccination will modify cases of chickenpox.
PPD	Mantoux Test for Tuberculosis
	<ol style="list-style-type: none">1. Return to office in 48-72 hours to have forearm checked.
Rotateq	Rotavirus vaccine
	<ol style="list-style-type: none">1. 1-3% incidence of mild, temporary diarrhea or vomiting within 7 days.